

Rx  
**SENSID**  
Sennosides Tablets USP 12 mg

**Composition:**

Each uncoated tablet contains:  
Sennosides USP.....12 mg  
Excipients.....q.s.  
Approved colour used.

**Pharmaceutical Form:**

Uncoated tablets

**Therapeutic indications:**

SENSID is indicated for, relief from occasional constipation.

**Posology and method of administration:**

**Posology:**

Take preferably at bedtime or as directed by the physician.  
Adults and children 12 years of age and over : 1 tablet once or twice daily.  
Children 6 to under 12 years of age : 1/2 tablet once or twice daily.  
Children under 6 years of age : ask a doctor before use.

**Method of administration:** For Oral administration.

**Contraindications:**

SENSID is contraindicated in patients with known hypersensitivity to sennosides or any the component of the formulation.

**Warnings and Precautions:**

Do not use, longer than one week, unless directed by the physician & when abdominal pain, nausea, or vomiting are present.  
Inform physician before use, if you have noticed a sudden change in bowel habits that persists over a period of 2 weeks.  
Consult a doctor or pharmacist before use, if you are taking any other drug.  
Take SENSID 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.  
Stop use and consult a physician, if you have rectal bleeding or fail to have a bowel movement after use of a laxative. These may be signs of a serious condition.

**Drug interactions:**

Hypokalaemia (resulting from long-term laxative abuse) potentiates the action of cardiac glycosides and interacts with antiarrhythmic medicinal products, with medicinal products which induce reversion to sinus rhythm (e.g. quinidine) and with medicinal products inducing QT-prolongation. Concomitant use with other medicinal products inducing hypokalaemia (e.g. diuretics, adrenocorticosteroids and liquorice root) may enhance electrolyte imbalance.

**Pregnancy and Lactation:**

SENSID should not be used, during pregnancy & lactation, without consulting the physician.

**Undesirable effects:**

Hypersensitivity reactions (pruritis, urticaria, local or generalized exanthema) may occur.  
SENSID may produce abdominal pain and spasm and passage of liquid stools, in particular in patients with irritable colon. However, these symptoms may also occur generally as a consequence of individual overdose. In such cases dose reduction is necessary.  
Chronic use may lead to disorders in water equilibrium and electrolyte metabolism and may result in albuminuria and haematuria. Furthermore, chronic use may cause pigmentation of the intestinal mucosa (pseudomelanosis coli).  
Yellow or red-brown (pH dependent) discolouration of urine by metabolites, which is not clinically significant, may occur during the treatment. The frequency is not known.

**Overdose:**

The major symptoms of overdose/abuse are griping pain and severe diarrhoea with consequent losses of fluid and electrolytes, which should be replaced. Diarrhoea may especially cause potassium depletion, which may lead to cardiac disorders and muscular asthenia, particularly where cardiac glycosides, diuretics, adrenocorticosteroids or liquorice root are being taken at the same time.  
Treatment should be supportive with generous amounts of fluid. Electrolytes, especially potassium, should be monitored. This is especially important in the elderly.  
Chronic ingested overdose of anthranoid containing medicinal products may lead to toxic hepatitis.

**Storage:** Store at a temperature below 30°C, in a dry place. Protect from light.  
Keep medicines out of reach of children.

**Presentation:** 100 Tablets in a jar along with package insert.

**MOH, Kuwait**

**Manufactured by:**

