

Rx

THIVITA

Thiamine Tablets BP 100 mg

For Oral Use Only

Composition:

Each Film Coated Tablet Contains:
Thiamine Hydrochloride BP.....100 mg
Excipients.....q.s.
Approved colour used.

Excipients : Dibasic calcium phosphate, Microcrystalline cellulose, Maize Starch, Lactose, Colloidal Silicon Dioxide, Polyvinylpyrrolidone K-30, Methylparaben, Propylparaben, Magnesium Stearate, Talcum, Sodium Starch Glycolate.

Pharmaceutical Form: Film Coated Tablets

Therapeutic indications: Treatment of thiamine deficiency.

Posology and method of administration:

Posology:

Adults, the Elderly and Children over three years:

Mild chronic deficiency: 50 mg daily
Severe deficiency: 100 mg three times daily.
Not recommended for children under three years.

Method of administration

For oral use.

Contraindications: Hypersensitivity to Thiamine Hydrochloride or to any of the excipients listed.

Warnings and Precautions:

Do not exceed recommended dosage.

This medicinal product contains lactose monohydrate. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.

Drug Interactions :

The thiamine antagonist's thiosemicarbazone and 5-fluorouracil can neutralise the effect of thiamine. Patients using any of these treatments may need their thiamine dose adjusted.

Thiamine could give false positive results for urobilinogen determination by the Ehrlich's reaction. High doses of thiamine may interfere with spectrophotometric assays of theophylline plasma concentration.

Pregnancy and Lactation:

This product is not intended for use in pregnant or lactating women.

Undesirable effects:

Gastrointestinal disorders:

Mild gastrointestinal events such as nausea, vomiting, diarrhoea, and abdominal pain have been reported. Frequency not known.

Immune system disorders:

Allergic and anaphylactic reactions, with symptoms of pruritus, urticaria, itching, hives, angioedema, abdominal pain, respiratory distress, tachycardia, palpitations, and shock have been reported in single cases. Frequency not known.

Overdose: In case of overdosage, treatment should be symptomatic.

Storage: Store below 30°C. Protect from light.

Keep the medicine out of reach of children.

Presentation: 100 tablets are packed in a jar along with pack insert.

MOH, Kuwait

Manufactured by:

