

ANIKOF-D

Syrup

Dextromethorphan Hydrobromide, Phenylephrine Hydrochloride,
Cetirizine Hydrochloride & Racementhol Syrup

Composition:

Each 5ml contains:	
Dextromethorphan Hydrobromide BP	10mg
Phenylephrine Hydrochloride BP	5mg
Cetirizine Hydrochloride BP	5mg
Racementhol BP	1.5mg
Excipients	qs
Flavoured syrup base	qs
Approved colour used.	

Pharmaceutical Form:

Oral Liquid.

Therapeutic indications:

ANIKOF-D is indicated for symptomatic relief of dry unproductive cough due to post nasal drip, allergic & vasomotor rhinitis, allergic pharyngitis, laryngitis & sinusitis, allergic cough following URTI and surgery.

Posology and method of administration:

Posology:

5-10 ml, 2-3 times a day.
Or as directed by the Physician.

Method of administration:

For oral administration.

SHAKE WELL BEFORE USE.

Contraindications:

ANIKOF-D is contraindicated in patients with pre-existing cardiac tachyarrhythmias.

Hypersensitivity to any of the ingredient of the formulation. And in conditions like Seizures, Hypersensitivity to murine proteins.

Warnings and Precautions:

ANIKOF-D, should not be administered to patients with chronic or persistent cough, such as occurs with asthma, or where cough is accompanied by excessive secretions, unless directed by a physician.

Administer ANIKOF-D with extreme caution to patients with hyperthyroidism, bradycardia, partial heart block, myocardial disease, prostatic hypertrophy, diabetes mellitus, increased IOP, or severe arteriosclerosis; trouble urinating due to enlarged prostate gland.

Pregnancy and Lactation:

ANIKOF-D be used only when the potential benefit of treatment to the mother exceeds any possible hazards to the developing foetus or suckling infant.

Drug interactions:

Dextromethorphan Hydrobromide: The concomitant use of a dextromethorphan-containing product and monoamine oxidase inhibitors can occasionally result in symptoms such as hyperpyrexia, hallucinations, gross excitation or coma.

Phenylephrine Hydrochloride: Concurrent use may enhance the pressor effects and induce tachycardia, especially in infants. Decrease phenylephrine's effect. Arrhythmias.

Cetirizine Hydrochloride: No clinically significant drug interactions have been found with theophylline at a low dose, azithromycin, pseudoephedrine, ketoconazole, or erythromycin.

Menthol: Menthol should be used only if clearly needed during pregnancy.

Undesirable effects:

Side effects attributed to dextromethorphan are uncommon; occasionally dizziness, nausea, vomiting, or gastro-intestinal disturbance may occur. Respiratory difficulty, Dysuria; urinary retention. Anorexia, flushing, increased salivation, urinary retention.

Overdose:

Overdose may include drowsiness, lethargy, nystagmus, ataxia, respiratory depression, nausea, vomiting, hyperactivity. Coma, hypotension, profuse sweating, sedation, sensation of fullness in head, severe hypertension, shock, short paroxysms of ventricular tachycardia, somnolence, tingling of extremities, ventricular extrasystoles, vomiting.

Storage:

Store below 30°C in dry place. Protect from light.

Keep the medicine out of reach of children.

Presentation:

100 ml bottle packed in a printed carton along with pack insert.

Registration No.: MCA/med 1919/07/25

Manufactured by :

