

ANIPEPTO

Bismuth Subsalicylate

Oral Suspension USP

Composition:

Each ml contains:

Bismuth subsalicylate USP	17.6 mg
Excipients	qs
Flavoured syrup Base	qs
Approved colour used.	

Pharmaceutical Form:

Oral Suspension.

Therapeutic indications:

ANIPEPTO is used as, upset stomach reliever and anti-diarrheal.

ANIPEPTO relieves:

- Diarrhea.
- Heartburn.
- Indigestion.
- Nausea.
- Upset stomach associated with these symptoms.

Posage and method of administration:

Posage:

Adults and children 12 years and over: 1 dose (30 ml or 2 TBSP) every 1/2 to 1 hour as needed.

Do not exceed 8 doses (240 ml or 16 TBSP) in 24 hours.

Use until diarrhea stops but not more than 2 days.

Children under 12 years: Consult your physician.

Drink plenty of clear fluids to help prevent dehydration caused by diarrhea.

Method of administration:

For oral administration.

Use measuring cup or tablespoon (TBSP).

SHAKE WELL BEFORE USE.

Contraindications:

Hypersensitivity to bismuth, aspirin, other salicylates.

Infectious diarrhea, high fever, von Willebrand disease, hemorrhage, ulcer or GI bleeding with black or bloody stool, hemophilia.

In paediatric patients, chicken pox or influenza (risk of Reye's syndrome).

Warnings and Precautions:

Bismuth subsalicylate may cause temporary and harmless darkening of tongue and/or black stools, generally reversible within several days after treatment stopped; stool darkening should not be confused with melena.

Bismuth absorbs x-rays and may interfere with x-ray diagnostic procedures of gastrointestinal tract.

Cases of neurotoxicity associated with excessive doses of various bismuth-containing products reported; effects have been reversible with discontinuation of bismuth therapy.

Not to be administered to children or teenagers recovering from flu-like symptoms or chickenpox; increased incidence of Reye's syndrome, associated with salicylate products.

If any of the below condition occurs, discontinue the use of Bismuth subsalicylate & consult the physician;

- diarrhea lasts more than 2 days.
- symptoms get worse or last more than 2 days.
- ringing in the ears or loss of hearing occurs.

Pregnancy and Lactation:

Consult the physician before use.

Drug interactions:

Do not take Bismuth subsalicylate with Methotrexate.

Bismuth subsalicylate may also interact with the following:

- Aspirin and aspirin-like medications.
- Medications for diabetes.
- Medications for gout.
- Medications that treat or prevent blood clots like warfarin.

Undesirable effects:

- Allergic reactions: skin rash, itching, hives, swelling of the face, lips, tongue, or throat.

- Bleeding: bloody or black, tar-like stools, vomiting blood or brown material that looks like coffee grounds, red or dark brown urine, red or purple spots on skin, unusual bruising or bleeding.

- Hearing loss, ringing in ears.

Side effects that usually do not require medical attention:

- Black stools.
- Constipation.
- Dark or black tongue.

Overdose:

Initiate general symptomatic and supportive measures in all cases of overdosages where necessary.

Storage:

Store below 30°C in a dry place. Protect from light.

Keep the medicine out of reach of children.

Presentation:

100 ml glass bottle packed in a printed carton along with pack insert.

Registration No.: MCA/med 1908/07/25

Manufactured by :



(INDIA) Private Limited
Pioneering Excellence. Redefining Quality.
www.uspharma.in