

BROMEXINA ADULTO

Bromhexine Hydrochloride Syrup 8mg/5ml

Composition:

Each 5 ml Contains
Bromhexine Hydrochloride BP 8mg
Excipients q.s.
Flavoured syrup base q.s.
Approved colour used

Dosage:

As directed by the physician

Storage: Store below 30°C in a dry place. Protect from light and moisture

Bromhexine 8 mg / 5 ml syrup**What is this medicine used for?**

Bromhexine is used for conditions where there is a large amount of thick mucus in the airways. Bromhexine belongs to a group of medicines called mucolytics. Bromhexine thins the mucus in the airways, thus making the mucus less sticky and also making it easier to remove the mucus.

How to take this medicine:

Adults and children over 2 years of age: Do not use for more than 14 days without medical advice.

AGE	RECOMMENDED DOSE FOR SYRUP
Adults and children 12 years and older	10 ml. to be taken 3 times a day
Children 6-12 years old	5 ml. to be taken 3 times a day
Children 2-5 years old	2.5 ml to be taken 3 times a day

Bromhexine can be taken with or without food. For the syrup, shake the bottle well before use.

If you miss a dose, take it as soon as you remember. However, if it is almost time for your next dose, skip the missed dose and continue with your next dose at the usual time. Do not take a double dose at once.

Before using this medicine

Medical advice should be sought before use if you have any of the following conditions: Symptoms indicative of pneumonia or other lung infections (e.g. difficulty breathing while resting, blood-stained mucus, fever with a temperature over 38°C, chest pain), a low immune system due to other health conditions (e.g. HIV) or medications (e.g. chemotherapy, immune system medication or lung conditions) such as chronic obstructive pulmonary disease or severe uncontrolled asthma. If you have gastric ulcers, liver or kidney problems This medicine should be avoided if you have a known history of allergy to bromhexine or any of the other listed ingredients of the product. Please consult your doctor or pharmacist before using this medicine if you are pregnant or breastfeeding or may become pregnant or breastfeeding during the course of medication.

Interactions with other medicines

Be sure to tell your doctor and/or pharmacist if you are taking, have recently taken or plan to take any other medicines, nutritional supplements or herbal products.

Tell your doctor or pharmacist if you are taking any antibiotics as bromhexine may increase the absorption of antibiotics.

Side effects

Occasional, mild side effects include: a feeling of fullness in the stomach (bloating), diarrhea, dizziness, headache, indigestion, nausea, sweating and skin rash.

When to seek medical advice

Tell your doctor and/or pharmacist immediately if you experience any allergic symptoms (e.g. skin lesions, rashes, itching, difficulty breathing or eye swelling) or serious side effects due to this medication.

KEEP THE MEDICINES OUT OF REACH OF CHILDREN.

SHAKE WELL BEFORE USE.

Manufactured by :



US Pharma[®]
(INDIA) Private Limited
Pioneering Excellence. Redefining Quality.
www.uspharma.in