

Appetite Stimulant

C-4 Forte Syrup

Cyproheptadine with Multivitamin Syrup

COMPOSITION:

Each 5 ml contains:

Cyproheptadine HCl	BP	2 mg
Vitamin B1	BP	2 mg
Riboflavin Sodium Phosphate	BP	2 mg
Pyridoxine Hydrochloride	BP	0.75 mg
D-Panthenol	BP	2.5 mg
Nicotinamide	BP	22.5 mg
Flavored syrup base		q.s.

Color: Approved color used.

PHARMACEUTICAL FORM: Syrup

PHARMACOTHERAPEUTIC CATEGORY: Antihistamine.

INDICATIONS:

Anorexia in children and adolescents, whether organic or psychological; post-infectious anorexia; anorexia due to deficiency; growth retardation; childhood anorexia; habitual vomiting (except when of organic origin); emotional instability accompanied by anorexic episodes; anorexia associated with constitutional thinness in adults.

CONTRAINDICATIONS:

Newborn or premature infants: This medication must not be used in newborns or premature infants. Breastfeeding mothers: Due to the increased risk posed by antihistamines to infants in general—and to newborns and premature infants in particular—antihistamine therapy is contraindicated in breastfeeding mothers.

DRUG INTERACTIONS:

MAO inhibitors prolong and intensify the anticholinergic effects of antihistamines. Antihistamines may have additive effects with alcohol and other CNS depressants, e.g., hypnotics, sedatives, tranquilizers, and anxiolytics.

SIDE EFFECTS AND SPECIAL PRECAUTIONS:

Always consult a doctor before giving cold or allergy medication to a child. Death may occur due to the misuse of cold and allergy medications in very young children. You should not use this medication if you are allergic to cyproheptadine or if you have narrow-angle glaucoma, a stomach ulcer or obstruction, an enlarged prostate or urinary problems, if you are having an asthma attack, or if you are elderly or have a debilitating illness. Before taking this medication, tell your doctor if you have glaucoma, a history of asthma, high blood pressure, heart disease, or an overactive thyroid. Cyproheptadine may impair your thinking or reactions. Be careful if you drive or do anything that requires alertness. Alcohol consumption may increase certain side effects of C4 FORTE.

SIDE EFFECTS:

Adverse reactions reported with the use of antihistamines are as follows:

Central nervous system

Sedation and drowsiness (often transient), dizziness, coordination disturbances, confusion, restlessness, excitement, nervousness, tremor, irritability, insomnia, paresthesia, neuritis, convulsions, euphoria, hallucinations, hysteria, fainting.

Skin Allergic manifestations such as skin rash and edema, excessive sweating, urticaria, photosensitivity.

EFFECTS ON THE ABILITY TO DRIVE AND USE MACHINES:

None anticipated.

LIST OF EXCIPIENTS:

Butyl hydroxybenzoate, Disodium edetate, Dilute hydrochloric acid, Ethyl hydroxybenzoate, Methyl hydroxybenzoate, Propyl hydroxybenzoate, Propylene glycol, Soluble orange oil flavoring, Sorbitol solution 70% (non-crystallizing), Purified water.

DOSAGE AND INSTRUCTIONS FOR USE:

ROUTE OF ADMINISTRATION: ORAL

Adults: Two 5 ml teaspoonfuls before breakfast, lunch, and dinner.

Children:

Infants: Two 2.5 ml teaspoonfuls per day, divided into two doses. Ages 7 to 12: One 5 ml teaspoonful before breakfast, lunch, and dinner, or one capsule before breakfast and one before dinner (2 days). Or as directed by your doctor.

CONSULT YOUR DOCTOR BEFORE TAKING THIS MEDICINE.

CHECK THE PRODUCT'S SHELF LIFE BEFORE USE. NOTICE:

STORAGE:

Store in a cool place at a temperature below 30°C.

Keep in the original packaging to protect from light and moisture.

PRESENTATION:

C4 FORTE Syrup is available in a 200 mL bottle.

Keep medicines out of the reach of children.

Manufactured by :

