

# Ciprofloxacin

## Ophthalmic Solution USP

**Composition:**

Ciprofloxacin Hydrochloride USP  
Ciprofloxacin.....0.3% w/v  
Benzalkonium Chloride USP....0.01% w/v  
(As a preservative)  
Sterile aqueous base.....q.s.

**Indication:**

Ciprofloxacin eye drops have a bactericidal action and are used to treat a variety of ophthalmic/otic infections. Avoid use in patients with a history of hypersensitivity to ciprofloxacin and its derivatives. Use with caution in the elderly population, as there is an increased risk of tendonitis and tendon rupture.

**Instructions:**

Apply ciprofloxacin eye drops according to your doctor's instructions. Do not exceed the recommended dosage. Consult your doctor if any side effects occur. Do not stop using ciprofloxacin eye drops before completing the prescribed course of treatment.

**Dosage:**

During the first two days of treatment, apply 1 drop every 2 hours during the day. On subsequent days, apply 1 drop every 4 hours during the day. Continue treatment until the infection has cleared. The duration of treatment should not exceed 10 days.

**Side Effects:**

Possible transient eye irritation, which should disappear after the first few applications. If it persists, consult your doctor.

**Contraindication:**

Patients with a history of hypersensitivity to ciprofloxacin.

**Warning:**

1. If irritation persists or worsens, discontinue treatment and consult your doctor.
2. Do not touch the dropper tip to any surface, as this may contaminate the solution.

**Storage:**

Store in a cool, dark place.

Keep the medication out of the reach of children.

**Presentation:** 10 mL bottle

**FOR EXTERNAL USE**

**NOT FOR INJECTION**

Manufactured in India:



**US Pharma**<sup>®</sup>  
(INDIA) Private Limited  
Pioneering Excellence. Redefining Quality.  
[www.uspharma.in](http://www.uspharma.in)