

Erythromycin

Stearate Tablets BP

Composition: Each film-coated tablet contains:

Erythromycin Stearate BP
Equivalent to Erythromycin 500 mg
Excipients q.s.
Color: Approved color used.

Uses:

Erythromycin is used to treat a wide variety of bacterial infections. It can also be used to prevent certain bacterial infections. Erythromycin is known as a macrolide antibiotic. It works by stopping the growth of bacteria.

This antibiotic treats or prevents bacterial infections only. It will not work for viral infections (such as the common cold or flu). Unnecessary use or misuse of any antibiotic can lead to decreased effectiveness.

Dosage and duration of treatment are based on your medical condition and response to treatment. In children, dosage is also based on age and weight.

For the best effect, take this antibiotic at evenly spaced intervals. To help you remember, take this medication at the same time(s) every day.

If you are using this medication to treat an infection, continue taking it until the prescribed amount is finished, even if symptoms disappear after a few days. Stopping the medication too soon may result in the infection returning.

Side effects: Nausea, vomiting, diarrhea, stomach pain/cramps, and loss of appetite may occur. Taking this medication with food may reduce these symptoms. If any of these effects persist or worsen, notify your doctor or pharmacist immediately. Remember that your doctor prescribed this medication because he or she determined that the benefit to you outweighs the risk of side effects. Many people using this medication do not experience serious side effects.

Precautions

Before taking erythromycin, tell your doctor or pharmacist if you are allergic to it, or to other macrolide antibiotics (such as azithromycin, clarithromycin), or if you have any other allergies.

This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details.

Before using this medication, tell your doctor or pharmacist your medical history, especially regarding: kidney disease, liver disease, or a certain type of muscle disease (myasthenia gravis).

Interactions

Drug interactions may change how your medications work or increase the risk of serious side effects. This document does not list all possible drug interactions. Keep a list of all products you use (including prescription and non-prescription medicines and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medication without your doctor's approval.

Overdose

If someone has overdosed and experiences severe symptoms, such as fainting or difficulty breathing, contact your doctor immediately.

Dosage: As directed by a physician.

Storage: Store below 30°C in a dry place; protect from light.

Keep medicines out of the reach of children.

Presentation: Blister pack of 10 tablets.

Manufactured by:

