

# HEMORRHOID

Hydrocortisone Acetate + Lidocaine + Zinc Oxide + Allantoin Ointment

## Treatment of external and internal hemorrhoids

### Composition:

Hydrocortisone acetate	BP	0.25% w/w
Lidocaine	BP	3.0% w/w
Zinc oxide	BP	5.0% w/w
Allantoin	BP	0.5% w/w
Ointment base		q.s.

**USES:** This medication contains an antibiotic that helps prevent or treat infection, an anesthetic that acts on the skin to relieve pain, and a steroid that reduces inflammation. It is used to treat hemorrhoids and other rectal conditions.

**HOW TO USE:** This medication is for rectal use only. For best results, use this medication as prescribed—usually once in the morning, once at night, and after each bowel movement. Do not use it more frequently or for a longer period than prescribed. Apply a small amount of the ointment to the affected rectal area as directed. If using inside the rectum, attach the rectal applicator to the tube. Carefully insert the applicator fully into the rectum. Gently squeeze the tube to dispense the medication while withdrawing the applicator. Wash the applicator and your hands after use.

**SIDE EFFECTS:** This medication may temporarily sting or burn for a minute or two when first applied. If this persists or becomes bothersome, inform your doctor. Tell your doctor if you develop a rash, or redness, itching, pain, or swelling around the rectal area. If you notice other effects not listed above, contact your doctor.

**PRECAUTIONS:** Inform your doctor if you have: other infections or diseases (especially colitis), intestinal obstruction, peritonitis, abscesses, fistulas, recent intestinal surgery, eye problems, or any allergies. Although highly unlikely, it is possible for this medication to be absorbed into the bloodstream. This could lead to undesirable consequences requiring additional treatment with corticosteroids. This is particularly relevant for children and individuals who have used the medication for an extended period, especially if they also have serious medical conditions such as severe infections, injuries, or surgeries. This precaution applies for up to one year after discontinuing use of this medication.

**Inform your doctor immediately if any of the following side effects occur:** vision problems, persistent headache, increased thirst or urination, weakness or unusual weight loss, or dizziness. Consult your doctor or pharmacist for further details, and inform them that you are taking (or have taken) this medication. Your dosage may need to be gradually reduced if treatment is no longer required and you have been taking multiple doses daily for an extended period. This medication should be used during pregnancy or breastfeeding only if clearly necessary. Discuss the risks and benefits with your doctor.

**DRUG INTERACTIONS:** Inform your doctor about any medication you may be taking. Do not start or stop taking any medication without medical or pharmaceutical approval.

**OVERDOSE:** If an overdose is suspected, contact the local poison control center or the emergency room immediately.

**MISSED DOSE:** If you miss a dose, use it as soon as you remember; do not use it if it is close to the time for your next dose. Instead, skip the missed dose and resume your usual dosing schedule. Do not "double up" the dose to make up for the missed one.

**Dosage and directions for use:** Apply locally 2 or 3 times a day as needed. For internal hemorrhoids, use the closed cannula. As directed by a physician.

Store at a temperature not exceeding 25°C.

For external use only.

Keep medicines out of the reach of children.

Presentation: One 20 g tube.

Manufactured by:



**US Pharma**<sup>®</sup>  
(INDIA) Private Limited  
Pioneering Excellence. Redefining Quality.  
[www.uspharma.in](http://www.uspharma.in)