

# Indomethacin Capsules

## BP 25 mg

**Composition:** Each hard gelatin capsule contains:

Indomethacin BP 25 mg

Excipients Q.S.

Approved color used in the capsule shell.

Indomethacin capsules belong to a group of medicines known as non-steroidal anti-inflammatory drugs (NSAIDs). These medicines are painkillers that reduce swelling.

Indomethacin capsules are used for:

- \* inflammatory joint diseases, such as rheumatoid arthritis, osteoarthritis, and degenerative hip disease
- \* pain and stiffness in the spine (ankylosing spondylitis)
- \* sprains and strains (acute muscle and bone disorders)
- \* lower back pain
- \* pain and swelling following muscle or bone surgery
- \* menstrual pain
- \* gout.

**Contraindications:**

- \* you are in the last three months of pregnancy
- \* you are allergic (hypersensitive) to indomethacin or to any of the ingredients in indomethacin capsules
- \* you have an allergy (hypersensitivity) to aspirin, ibuprofen, or other NSAIDs, or have developed signs of asthma (wheezing), a runny nose, swelling of the face, lips, tongue, or throat, or a skin rash with irregular pale or red patches accompanied by intense itching, when taking these medicines
- \* you have severe liver, kidney, or heart problems
- \* you have nasal obstruction (nasal polyps).

**Precautions:** Before taking indomethacin, tell your doctor or pharmacist if you are allergic to it; or to aspirin or other NSAIDs (such as ibuprofen, naproxen, celecoxib); or if you have other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details.

**Interactions:** Drug interactions may alter the way your medications work or increase the risk of serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription and over-the-counter medications, and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medication without your doctor's approval.

**Pregnancy and Breastfeeding:** If you are in the last three months of pregnancy, you should not take Indometacin capsules. You should only take Indometacin capsules during the first six months of pregnancy or while breastfeeding under medical supervision. Always consult your doctor before taking any medication.

**Side Effects:** This medication may raise your blood pressure. Check your blood pressure regularly and inform your doctor if the readings are high.

**Dosage:** As directed by your doctor.

**Storage:** Store below 30°C in a cool, dry place. Protect from light.

**Keep out of reach of children.**

**Presentation:** Blister pack of 10 capsules.

**Manufactured by:**



**US Pharma**<sup>®</sup>  
(INDIA) Private Limited  
Pioneering Excellence. Redefining Quality.  
[www.uspharma.in](http://www.uspharma.in)