

Mebendazole Tablets USP 100mg

Linozole

Composition:

Each uncoated tablet contains:

Mebendazole USP 100 mg

Excipients qs

Approved colour used.

Dosage:

As directed by the physician.

Indications:

Treatment of infestations by *Ascaris lumbricoides*, *Enterobius vermicularis*; *Trichiura*; *Ancylostoma duodenale* and *Trichinella spiralis*.

Side Effects: Generally well tolerated. Occasionally transient gastrointestinal disturbances, headache and rarely, allergic reactions and reversible neutropenia.

Contraindications: History of hypersensitivity to mebendazole and in the first trimester of pregnancy.

Notes and Precautions:

- (1) Administer preferably between meals.
- (2) In generalized infections, mainly caused by *Ancylostoma* and *Enterobius*, it is advisable repeat a new treatment cycle after 2-4 weeks.
- (3) In trichinosis, extend treatment for 5 days, combining, if necessary, prednisolone for relief of allergic and inflammatory symptoms.
- (4) Since its safety has not been well established, avoid use in children under 2 years of age and during pregnancy (teratogenic and embiotoxic effects).

Presentation: Blister of 6 tablets.

Storage: Store below 30°C in a dry place, protect from light.

Do not use during pregnancy.

Keep the medicines out of reach of children.

Manufactured by:



US Pharma[®]
(INDIA) Private Limited

Pioneering Excellence. Redefining Quality.
www.uspharma.in