

Loperamide Hydrochloride Tablets USP 2 mg

Composition: Each uncoated tablet contains:
Loperamide Hydrochloride USP 2 mg
Excipients q.s.

Description

Loperamide HCl 2 mg Tablets are a highly effective anti-diarrheal medication. They are used to relieve the symptoms of diarrhea that begins suddenly and lasts for a few days. They are also used to treat long-term diarrhea associated with other medical conditions. The effects of this medication can be observed within 1 to 3 hours after administration. Loperamide is typically administered once at the start of treatment, and then after each loose stool (diarrhea), up to four times a day, until the diarrhea improves. It should not be taken more frequently than every 3 hours or for more than 2 days.

Warnings and Precautions

Pregnancy: The use of this medication is not recommended for pregnant women unless the potential benefits outweigh the associated risks. Consult your doctor before taking this medication.

Breastfeeding: The use of this medication is not recommended while breastfeeding an infant. If the use of this medication is absolutely necessary, breastfeeding should be discontinued. Consult your doctor before taking this medication.

Hypersensitivity: The use of this medication may cause allergic skin reactions in some patients. Some cases of severe allergic conditions, such as Stevens-Johnson Syndrome or toxic epidermal necrolysis, have been reported with the use of this medication.

Cardiac Conditions: The use of this medication in large quantities may cause severe adverse effects on the heart and blood vessels. Non-prescribed use of this medication should be avoided.

Pediatric Use: This medication should be used with caution in patients under 6 years of age. Use in children under 2 years of age is not recommended.

Constipation: The use of this medication is not recommended for patients suffering from intestinal obstruction or constipation.

Infectious Diarrhea: The use of this medication is not recommended for infectious diarrhea characterized by the presence of pus or blood in the stool, high fever, etc. Acute diarrhea and dysentery caused by bacteria should not be treated with this medication.

General Instructions

Take Loperamide 2 mg tablets as directed by your doctor. This medication may be taken with or without food. Do not take more than the prescribed dose or use it for longer than recommended. It is recommended to drink sufficient fluids while taking this medication. Contact your doctor if symptoms do not improve within 48 hours of taking this medication.

Side Effects

Dizziness, drowsiness, tiredness, or constipation may occur. If any of these effects persist or worsen, contact your doctor immediately.

If your doctor has prescribed this medication, remember that he or she has judged that the benefit to you outweighs the risk of side effects. Many people who use this medication do not experience serious side effects.

Dosage and Administration: Oral: As directed by a physician.

Storage: Store below 30°C in a cool, dry place. Protect from light.

Presentation: Each blister contains 10 tablets.

KEEP MEDICINES OUT OF THE REACH OF CHILDREN.

Manufactured by:

