

Lysine Acetylsalicylate

For solution for injection

Composition:

Each glass vial contains:

Lysine acetylsalicylate	0.9 g
Glycine BP	0.1 g

Indications:

Moderate to severe pain of various origins, rheumatic pain, hyperthermia.

Contraindications:

Gastroduodenal ulcer, history of hypersensitivity to salicylates; any bleeding disorders (hereditary or acquired); any risk of hemorrhage; last trimester of pregnancy. Use is not recommended in cases of gout or menorrhagia.

Side effects:

The side effects are the same as those of acetylsalicylic acid.

Precautions:

Use with caution in cases of: concomitant anticoagulant therapy, asthma, renal insufficiency, the first two trimesters of pregnancy, and IUD use.

Drug interactions:

Vitamin K antagonists and heparin, sulfonyleureas, methotrexate, NSAIDs, uricosuric agents.

Dosage and administration:

Adults: 1 or 2 vials per injection;
for severe pain: 2 vials in a single IV injection.

The usual 24-hour dosage is 1 to 4 vials.

In some cases, the daily dose may be increased to 8 vials.

Children: 10–25 mg/kg/day. For children aged 6–10 years: 1/2 vial per day.

Over 11 years: 1–2 vials per day.

Storage: Store below 25°C.

Keep all medicines out of the reach of children.

Manufactured by :



US Pharma[®]
(INDIA) Private Limited
Pioneering Excellence. Redefining Quality.
www.uspharma.in