

MEXOGRA-100



Sildenafil Tablets BP 100 mg

Composition:

Each film coated tablet contains:

Sildenafil Citrate BP

Eqvt. to Sildenafil.....100 mg

Excipients.....q.s.

Approved colour used

Pharmaceutical form : Film coated tablet.

Indications: Management of erectile dysfunction.

Side effects: The most common are headache and flushing; They can also arise visual disturbances, dizziness and conjunctival congestion, muscle pain, rash, urinary infection and priapism; also were reported, palpitations and severe cardiac all traces including sudden death.

Contraindications: Patients with severe hepatic insufficiency, antecedents of bleeding, peptic ulcer active, arterial hypotension, stroke, myocardial infarction, arrhythmia, unstable angina, cardiac insufficiency and alterations of the retina. Patients treated with organic nitrates, patients taking ritonavir or saquinavir.

Notes and Precautions:

- (1) Reduce the dose in patients with renal or hepatic insufficiency.
- (2) Use with caution in patients with hematological alterations that can precaution in donets with hematological alterations that may predispose to priapism.
- (3) Patients with impaired vision or dizziness should not drive or operate heavy machinery.
- (4) SILDENAFIL TABLETS BP 100 mg may potentiate the hypotensive effects of organic nitrates.
- (5) drugs such as cimetidine, erythromycin, itraconazole, ketoconazole and HIV protease inhibitors may reduce the clearance of SILDENAFIL TABLETS BP 100 mg in need of a reduction in the dose.
- (6) The grape juice should be avoided because it may increase the plasma levels of SILDENAFIL TABLETS BP 100 mg.
- (7) It should not be used in men whose sexual activity and inadvisable due to underlying cardiovascular disease.
- (8) Patients should be advised that having taken SILDENAFIL TABLETS BP 100 mg in the last 24 hours, should not take infirm medication without consulting the doctor.
- (9) It should be consulted the doctor, when the reversal of erections persists beyond 4 hours so as to avoid fibrosis of the corpora cavernosa and therefore a permanent erectile dysfunction.
- (10) No adequate cause an erection in the absence of sexual stimulation.
- (11) Do not have an aphrodisiac effect.

Dosage: As directed by the Physician.

Storage: Store below 30°C in dry place. Protect from light.

Keep the medicine out of reach of children.

Method of Administration : For Oral Administration.

Presentation : A carton containing 2 blisters of 4 tablets along with pack insert

Manufactured by:



US Pharma[®]
(INDIA) Private Limited
Pioneering Excellence. Redefining Quality.
www.uspharma.in