

MUSCLE PLUS

Comprimidos

Composition: Each uncoated tablet contains:

Diclofenac Potassium	BP	50 mg
Chlorzoxazone	USP	250 mg
Paracetamol	BP	500 mg
Excipients		q.s.

Color: Approved color used.

Indication: Chlorzoxazone + Diclofenac + Paracetamol Tablets are indicated for the treatment of muscle pain, fever, headache, febrile conditions, joint pain, relief of pain associated with osteoarthritis of the

knees and hands, earache, muscle spasms, pain and stiffness due to muscle strains and sprains, toothache, and other conditions. Chlorzoxazone + Diclofenac + Paracetamol Tablets may also be used for purposes not mentioned in this medication guide.

Chlorzoxazone + Diclofenac + Paracetamol Tablets contain the following active ingredients: Chlorzoxazone, Diclofenac, and Paracetamol.

Mechanism of Action and Pharmacology: Chlorzoxazone + Diclofenac + Paracetamol Tablets improve the patient's condition by performing the following functions:
Depressing muscle reflexes.

Blocking the action of the cyclooxygenase enzyme, which is necessary for the biosynthesis of prostaglandins.

Increasing the pain threshold and increasing blood flow through the skin, heat loss, and sweating.

Side Effects:

The following is a list of possible side effects that may occur with medications containing Chlorzoxazone + Diclofenac + Paracetamol. This is not a comprehensive list.

These side effects are possible,

but do not always occur. Some side effects may be rare but serious.

Consult your doctor if you observe any of the following side effects, especially if they do not go away. Allergic reactions

Coughing up blood or vomiting

Gastrointestinal bleeding

Drowsiness

Dizziness

Lightheadedness

Malaise

Skin rashes

Angioedema

Anaphylactic reactions

Precautions:

Before using Chlorzoxazone + Diclofenac + Paracetamol Tablets, inform your doctor about your current list of medications, over-the-counter products (e.g., vitamins, herbal supplements, etc.),

allergies, pre-existing diseases, and current health conditions (e.g., pregnancy, upcoming surgery, etc.).

Certain health conditions may make you more susceptible to the side effects of this medication. Take it exactly as directed by your doctor or follow the instructions printed on the package insert. The dosage is based on your medical condition. Inform your doctor if your condition persists or worsens. Important

advisory points are listed below:

Avoid contact of this medication with the eyes.

Avoid consuming alcohol or taking other sedative medications.

Avoid driving or operating machinery until you know how this medication affects you.

Avoid using this medication if you are allergic to paracetamol.

Blood pressure should be monitored closely during treatment.

Concomitant use with other sedatives.

Do not take paracetamol if you consume alcoholic beverages on a daily basis.

Elderly patients

Other allergies

Patients taking diuretics and angiotensin-converting enzyme (ACE) inhibitors

Drug Interactions:

If you take other medications or non-prescription products at the same time, the effects of the Chlorzoxazone + Diclofenac + Paracetamol Tablet may change. This may increase the risk of side effects or cause the medication to not work properly. Inform your doctor about all drugs, vitamins, and herbal supplements you are taking so that the doctor can help you prevent or manage drug interactions.

Contraindications:

Hypersensitivity to the Chlorzoxazone + Diclofenac + Paracetamol Tablet is a contraindication. Furthermore, the Chlorzoxazone + Diclofenac + Paracetamol Tablet should not be used if you have the following conditions:

- Hepatic insufficiency
- History of asthma, urticaria, or other allergic-type reactions after taking aspirin or other non-steroidal anti-inflammatory drugs (NSAIDs)
- Hypersensitivity
- Hypersensitivity to this drug
- Treatment of perioperative pain in the setting of coronary artery bypass graft (CABG) surgery
- Avoid during pregnancy and breastfeeding

Overdose:

Do not take more than the prescribed dose. Taking more medication will not improve your symptoms; instead, it may cause poisoning or serious side effects. If you suspect that you or anyone else has taken an overdose of

Chlorzoxazone + Diclofenac + Paracetamol Tablet, go to the nearest hospital emergency room or medical center. Bring the medicine box, container, or label with you to provide the doctors with the necessary information. Do not give your medication to other people, even if you know they have the same condition or it appears they may have similar conditions. This may lead to an overdose.

Dosage: As directed by a physician.

Keep medicines out of the reach of children.

Storage: Store below 30°C in a dry place; protect from light.

Presentation: Blister pack of 10 tablets.

Manufactured by:

