

PHENICLOR Chloramphenicol Eye Drops BP 0.5%

Composition:

Chloramphenicol BP	0.5% w/v
Phenyl Mercuric Nitrate BP (As Preservative)	0.02% v/v
Sterile Aqueous base	q.s.

WHAT CHLORAMPHENICOL EYE DROP B.P. 0.5 % W/V ARE AND WHAT THEY ARE USED FOR:

Chloramphenicol Eye Drop B.P. 0.5 % w/v contains a medicine called chloramphenicol. This belongs to a group of medicines called antibiotics. It works by stopping the growth of bacteria, allowing your body to fight off the infection.

Chloramphenicol Eye Drop B.P. 0.5 % w/v are used to treat minor infections of the eye or eyelid caused by bacteria (bacterial conjunctivitis). Signs include sore, red or inflamed eyes, sticky discharge or crust on the lids.

BEFORE USE:

Do not use this medicine and tell your doctor if: You are allergic (hypersensitive) to chloramphenicol or any of the other ingredients used. Signs of an allergic reaction include: a rash, swallowing or breathing problems, swelling of your lips, face, throat, tongue and worsening of redness, itching or swelling of the eye or eyelid. Do not use this medicine if the above apply to you. If you are not sure, talk to your doctor or pharmacist before using Chloramphenicol Eye Drop B.P. 0.5 % w/v.

Warnings and Precautions: You have problems with your eyesight, You have severe pain within the eye. Your eyes are sensitive to light, You have eye inflammation and a rash on your face or scalp, Your eye looks cloudy, The pupil of your eye looks unusual, You think that you have a foreign object in your eye, You have recently had conjunctivitis, You have glaucoma.

Pregnancy and breast-feeding: Talk to your doctor before using this medicine if you are pregnant, might become pregnant, or think you may be pregnant.

you should not breast-feed if you are taking Chloramphenicol Eye Drop. This is because small amounts may pass into mothers' milk. If you are breast-feeding or planning to breast-feed, talk to your doctor or pharmacist before taking or using any medicine.

Driving and using machines: You may have blurred eyesight straight after using this medicine. If this happens, do not drive or use any tools or machines until you can see clearly.

Chloramphenicol Eye Drop contains Boric acid: Do not give to a child less than 2 years old as this medicine contains boron and may impair fertility in the future.

HOW TO TAKE:

Always use Chloramphenicol Eye Drop exactly as your doctor or pharmacist has told you. You should check with your doctor or pharmacist if you are not sure.

Sing this medicine: Use the eye drops during waking hours only, Do not use for more than 5 days without talking to your doctor.

How to use this medicine: Wash your hands, Remove the cap from the bottle, Tilt your head back, Squeeze one drop inside the lower lid without touching your eye with the bottle, Close your eye, Wipe away any excess liquid from the eye with a clean tissue, Always put the cap back on the bottle as soon as you have used it.

How much to use

Adults, Elderly and Children over 2 years of age: One drop in the affected eye every two hours for the first 2 days (48 hours), After the first 2 days, one drop in the affected eye every four hours, Talk to your doctor or pharmacist if your symptoms do not improve within 2 days or if your eyes get worse, Keep using the eye drops for 5 days even if your symptoms improve.

If you forget to use Chloramphenicol Eye Drop: If you forget a dose, use your drops as soon as you remember. However, if it is nearly time for the next dose, skip the missed dose. Do not use a double dose to make up for a forgotten dose.

If you stop using Chloramphenicol Eye Drop : Keep using the eye drops for 5 days even if your symptoms improve. Do not stop using Chloramphenicol Eye Drop just because your eye(s) feel better. If you stop, your infection may come back. If you have any further questions on the use of this product, ask your doctor or pharmacist

POSSIBLE SIDE EFFECTS:

Like all medicines, Chloramphenicol Eye Drop can cause side effects, although not everybody gets them. Stop using Chloramphenicol Eye Drop and see a doctor as soon as possible if: You get any kind of skin problem, such as a rash or itching around your eyes or they become more red, You bruise more easily than usual or you get infections more easily or frequently than usual. You may also feel tired or weak; You notice any swelling of your face, lips or tongue, itching or rash with swelling of skin. Tell your doctor or pharmacist if any of the following side effects get serious or lasts longer than a few days: Stinging or burning in your eyes. This should only last for a short time and occurs immediately after using the eye drops.

HOW TO STORE:

Store in a cool place, Protect from light.

PACK PRESENTATION:

10 ml eye drop plastic bottle with mono carton and Insert.

Manufactured in India:



US Pharma[®]
(INDIA) Private Limited
Pioneering Excellence. Redefining Quality.
www.uspharma.in