

Salbutamol

Oral Solution BP

Composition: Each 5 ml contains:

Salbutamol Sulfate BP

equivalent to Salbutamol 2 mg

flavored syrup base q.s.

Color: Approved color used.

Contraindications: Salbutamol should not be taken concomitantly with beta-blocking agents. Hyperthyroidism and heart disease. Safety during pregnancy and lactation has not been established. Salbutamol is not indicated for the prevention of premature labor associated with toxemia of pregnancy or antepartum hemorrhage during the second and third trimesters of pregnancy.

Interactions: The adverse effects of high doses of salbutamol may be exacerbated by the concomitant administration of high doses of corticosteroids. The effects of salbutamol are antagonized by propranolol and other beta-blockers and potentiated by aminophylline or other xanthines. An increased risk of arrhythmias may occur if the patient is receiving cardiac glycosides, quinidine, or tricyclic antidepressants; interactions with alpha- and beta-blockers may also occur.

Side effects: GI disturbances such as nausea, vomiting, epigastric or retrosternal burning, and diarrhea. Other side effects may include palpitations, delayed skeletal muscle tremor (especially in the flank), and muscle cramps. Side effects can be managed by reducing the dosage.

Dosage:

Children aged 2 to 5 years: 5 ml, 3 times a day.

Children aged 6 to 14 years: 5 ml, 3 or 4 times a day.

Adults and children over 14 years: 5 to 10 ml, 3 or 4 times a day.

As directed by a physician.

Storage: Store below 30°C in a dry place; protect from light.

Keep medicines out of the reach of children.

SHAKE BEFORE USE.

Presentation: 100 ml bottle.

Manufactured by :

