

Kaolin, Neomycin Sulfate, and Pectin Suspension

Trans **KAOLIN** **SUSPENSION**

Composition:

Each 100 ml contains:

Kaolin	BP	17.50 g
Pectin	USP	440 mg
Neomycin Sulfate	BP	1.015 g
Flavored syrup base		q.s.

Color: Approved color used.

Dosage:

Children under 2 years: $\frac{1}{2}$ –1 teaspoonful 4 to 6 times daily

Children over 2 years: 2–3 teaspoonfuls 4 to 6 times daily

Adults: 2–3 teaspoonfuls three times daily, or as directed by a physician.

Indications: TRANSKAOLIN is indicated for the symptomatic treatment of mild to moderate acute diarrhea. It is also recommended for the symptomatic treatment of chronic diarrhea pending determination of the etiology.

Contraindications: This preparation should be avoided by patients hypersensitive to kaolin or pectin, as well as by patients suffering from diarrhea secondary to pseudomembranous enterocolitis or toxigenic bacteria.

Precautions: This preparation should be avoided in cases of suspected parasitic diarrhea, as it may make identifying the causative parasite more difficult. Furthermore, it should be avoided in acute dysentery characterized by bloody stools and high fever; in such cases, antibiotic therapy may be required. Caution is advised in patients suffering from dehydration, as adsorbent antidiarrheals may mask its severity. Consult a physician if diarrhea is not controlled within 48 hours or if fever develops.

Pregnancy and lactation: This medication is considered safe, as it is not absorbed into the body and is unlikely to cause problems.

Side effects: No serious side effects have been reported with the use of TRANSKAOLIN. However... Mild and transient constipation may occur in some patients.

Drug interactions: When taken together, TRANSKAOLIN may reduce the absorption of aspirin, chloroquine, hydroxychloroquine, phenothiazines, quinidine, digoxin, lincomycin, clindamycin, and tetracyclines. It is recommended that TRANSKAOLIN be administered at least 2 hours before, or 3 to 4 hours after, lincomycin and clindamycin. Patients receiving digoxin should be carefully monitored for evidence of altered effects. If you are taking any other medication, do not take it within 2 to 3 hours of taking TRANSKAOLIN. Concomitant administration may prevent the other medication from being absorbed by your body.

Storage: Store below 30°C in a dry place; protect from light.

Keep medicines out of the reach of children.

SHAKE BEFORE USE.

Presentation: 100 ml bottle.

Manufactured by :



US Pharma[®]
(INDIA) Private Limited
Pioneering Excellence. Redefining Quality.
www.uspharma.in