

Tramadol Hydrochloride Injection 100mg/2ml

Para uso IM/IV

COMPOSITION

Each ml contains

Tramadol Hydrochloride BP 50 mg
Water for Injection BP q.s.

GENERAL INFORMATION

It is centrally acting synthetic analgesic compound that is not derived from natural sources nor is chemically related to opiates.

CATEGORY

Opioid receptor agonist, noradrenaline reuptake inhibitor; analgesic.

INDICATIONS

Moderate to severe acute or chronic pain & in painful diagnostic measures & surgery. Arthralgia, musculoskeletal pain, colics, low back pain, pain associated with fractures, dislocation, contusion and osteoarthritis

DOSAGE AND ADMINISTRATION

Adult: Inj: 100 mg I.M. or S.C. or IV. to be injected slowly or diluted in solution for infusion. Maint: 50-100 mg every 4-6 hrs. Max 600 mg daily

Children: Not recommended for children below 14 yrs.

CONTRAINDICATIONS

Hypnotics, alcohol and psychotropic intoxication. Hypersensitivity.

SPECIAL PRECAUTION

Alcohol. Not to drive and operate machinery.

Tramadol should be used with caution in patients with pre-existing kidney disease, liver disease, seizure disorder, lungs disease, history of drug dependency or have any allergy. It should be used with caution during pregnancy or lactation.

ADVERSE EFFECTS

Nausea, vomiting and dizziness

INTERACTIONS

Carbamazepine: A significant increase in Tramadol metabolism.

Hypnotics, Tranquilizers and other CNS depressants. Sedative effects potentiated.

MAOIs: Fatal interaction-occurs

Opioid analgesics like Buprenorphine, Butorphanol and Pentazocine, May precipitate withdrawal symptoms

Tranquilizers Analgesia potentiated.

STORAGE

Storage : Store below 25°C. Protect from light

Keep out of the reach of children.

PRESENTATION

10 ampoules of 2ml enclosed in a plastic tray and each tray packed in a printed box.

Manufactured in India:



US Pharma[®]
(INDIA) Private Limited
Pioneering Excellence. Redefining Quality.
www.uspharma.in