

VFLIX-S®

Composition:

Each 5 ml contains:

Diphenhydramine Hydrochloride	BP	14.08 mg
Ammonium Chloride	BP	138.00 mg
Sodium Citrate	BP	57.03 mg
Menthol	USP	0.90 mg
Flavoured syrup base		q.s.

Colour: Approved colour used.

Indications : VFLIX-S syrup is used for the treatment, control, prevention, & improvement of the following diseases, conditions and symptoms: Relief of runny nose, sneezing, itchy, watery eyes, itchy nose, itchy throat due to hay fever or allergy. Relief of runny nose and sneezing due to common cold, Acidity, Back pain, Pain in shoulder joint, Appendicitis, Cough, Patients with metabolic alkalosis, Patients with hypochloremic states, Pain in arthritis, Pain in tendons, Pain in muscle strains or sprains, Bruising, Cramping, Gastric problems and Kidney stones.

Dosage: Children over 2 years: 1 teaspoonful (5 ml) every 4 hours.

Adults: 2 teaspoonful (10 ml) every 4 hours.

Or as directed by the physician.

Contraindications: VFLIX-S Syrup is a contraindicated in hyper sensitivity to Diphenhydramine Hydrochloride, Ammonium Chloride, Sodium Citrate and Menthol. VFLIX-S Syrup should not be used if you have the following conditions:

Acute dehydration, Anuria, Azotemia, Concomitant use with other products containing diphenhydramine, Heat cramps, Hyperkalemia, Hypersensitivity, Impaired renal function with oliguria, Metabolic alkalosis with loss of sodium bicarbonate and Seizures.

Precautions : Before using VFLIX-S Syrup inform your doctor about your current list of medications, over the counter products (e.g. vitamins, herbal supplements, etc.), allergies, pre-existing diseases, and current health conditions (e.g. pregnancy, upcoming surgery, etc.). Consult a doctor if you have cardiac failure, hypertension, impaired renal function or peripheral and pulmonary edema. Inform your doctor if you have previous kidney or liver failure. Patients with high CO₂ and buffer base. Possible ammonia toxicity. Pregnant, planning to become pregnant or breast feeding, Presence of a breathing problem such as emphysema or chronic bronchitis.

Drug Interactions : VFLIX-S Syrup may interact with the following drugs and products:

Alcohol, Alginic acid, Aluminiumhydroxide, Anisindione, Aspirin, Calcium carbonate, Dicumarol, Methadone, Other products containing diphenhydramine and Sedatives

Pregnancy and lactation: Pregnancy: avoid. Potential risk to the foetus has been reported. Consult your doctor before use, or if you are planning to fall pregnant

Breastfeeding: avoid. It is unknown how this medication may affect your baby. Consult your doctor before use.

Side-effects : The following is a list of possible side-effects that may occur from all constituting ingredients of VFLIX-S Syrup.

This is not a comprehensive list. These side-effects are possible, but do not always occur. Some of the side-effects may be rare but serious. Consult your doctor if you observe any of the following side-effects, especially if they do not go away. Injection site reactions, Rash, Fever, Irregular breathing, Bradycardia, Progressive drowsiness, Mental confusion, twitching, Convulsions and Coma. VFLIX-S Syrup may also cause side-effects not listed here.

Overdosage : Do not use more than prescribed dos. Taking more medication will not improve your symptoms; rather they may cause poisoning or serious side-effects. If you suspect that you or anyone else who may have overdosed of VFLIX-S Syrup, please go to the emergency department of the closest hospital or nursing home. Bring a medicine box, container, or label with you to help doctors with necessary information.

Storage: Store below 30°C in a cool & dry place, protect from light.

SHAKE WELL BEFORE USE

Keep medicines out of reach of children.

Presentation : Bottle of 100 ml

© Registered Trade Mark.

Manufactured by :

