

# Vita SCORPION™

## 75 mg / 3 ml

### Diclofenac Sodium Injection

Anti-rheumatic, Anti-inflammatory, Analgesic.  
I.M. (Intrajlut.) / I.V. Infusion

#### Composition:

Each 3 ml contains:

Diclofenac Sodium BP.....75 mg

Benzyl Alcohol BP..... 4%w/v

(As preservative)

Water for Injection BP.....q.s.

#### Indications:

The injection contains Diclofenac Sodium which is used to alleviate all degrees of pain and inflammation in a wide range of conditions, including: a) conditions arthritic conditions, such as rheumatoid arthritis, osteoarthritis, juvenile chronic arthritis, spondylitis ankylosing, acute gout. b) Acute musculoskeletal diseases, such as peri-arthritis (e.g. frozen shoulder), tendonitis, tenosynovitis, bursitis. w) Other painful conditions resulting from trauma including fracture, low back pain, sprains, strains, dislocations, pain control and inflammation in surgeries orthopedic, dental and other minor pain, post-operative pain, colic pain kidney etc.

#### Contraindications:

Diclofenac Sodium contraindicated in individuals with hypersensitivity to damn it. History of asthma, hives, or allergic-type reactions after taking aspirin or other NSAIDs. Severe, sometimes fatal, anaphylactic reactions to NSAIDs have been reported in such patients are contraindicated.

#### Attention:

Diclofenac Sodium cannot be expected to replace corticosteroids or corticosteroid insufficiency. Abrupt discontinuation of corticosteroids may lead to exacerbation of the disease. Patients on prolonged therapy with corticosteroid therapy should be tapered slowly if the decision is made to discontinue corticosteroids and the patient should be observed closely for any evidence of adverse effects, including adrenal insufficiency and exacerbation of arthritis symptoms.

#### Side effects:

- \* upset stomach, mild heartburn or upset stomach, diarrhea, bloating from constipation, gases;
- \* dizziness, headache, nervousness;
- \* itchy skin or rash;
- \* blurry vision; or.
- \* ringing in the ears.

#### Dosage:

As directed by the doctor.

#### Excess dosage:

Symptoms following acute NSAID overdose have typically been limited to lethargy, drowsiness, nausea, vomiting and epigastric pain, which have been generally reversible with supportive treatment. Gastrointestinal bleeding occurred.

#### Drug interactions:

Drug interactions can change how your medications work or increase the risk of serious side effects. Keep a list of all products you use (including prescription/nonprescription medications) prescription and herbal medicines) and share them with your doctor. Do not start, stop or change the dosage of any medicine without your doctor's approval.

#### Storage Condition:

Store in a cool, dry & dark place.

#### Keep the medicine out of reach of children.

**Presentation:** 10 X 3ml Ampoules in each mono box.

Manufactured in India:

